

# Number of revolutions of a particle around a black hole: Is it infinite or finite?

Yuri V. Pavlov<sup>1,2</sup> · Oleg B. Zaslavskii<sup>2,3</sup> 

Received: 12 July 2017 / Accepted: 14 December 2017 / Published online: 28 December 2017  
© Springer Science+Business Media, LLC, part of Springer Nature 2017

**Abstract** We consider a particle falling into a rotating black hole. Such a particle makes an infinite number of revolutions  $n$  from the viewpoint of a remote observer who uses the Boyer–Lindquist type of coordinates. We examine the behavior of  $n$  when it is measured with respect to a local reference frame that also rotates due to dragging effect of spacetime. The crucial point consists here in the observation that for a nonextremal black hole, the leading contributions to  $n$  from a particle itself and the reference frame have the same form being in fact universal, so that divergences mutually cancel. As a result, the relative number of revolutions turns out to be finite. For the extremal black hole this is not so,  $n$  can be infinite. Different choices of the local reference frame are considered, the results turn out to be the same qualitatively. For illustration, we discuss two explicit examples—rotation in the flat spacetime and in the Kerr metric.

**Keywords** Black holes · Kerr metric · Rotating Frames

---

✉ Oleg B. Zaslavskii  
zaslav@ukr.net

Yuri V. Pavlov  
yuri.pavlov@mail.ru

<sup>1</sup> Institute of Problems in Mechanical Engineering, Russian Academy of Sciences, 61 Bol'shoy pr., St. Petersburg, Russia 199178

<sup>2</sup> N.I. Lobachevsky Institute of Mathematics and Mechanics, Kazan Federal University, 18 Kremlyovskaya St., Kazan 420008, Russia

<sup>3</sup> Department of Physics and Technology, Kharkov V.N. Karazin National University, 4 Svoboda Square, Kharkiv 61022, Ukraine